

MLL Monoclonal Antibody

Catalog No :	YM0444			
Reactivity :	Human			
Applications :	WB;IHC;IF;ELISA			
Target :	MLL			
Fields	>> Lysine degradation:>> Metabolic pathways:>> Cushing			
Fields.	syndrome;>>Transcriptional misregulation in cancer			
Gene Name :	MLL			
Protein Name :	Histone-lysine N-methyltransferase MLL			
Human Gene Id :	4297			
Human Swiss Prot No :	Q03164			
Mouse Swiss Prot	P55200			
Immunogen :	Purified recombinant fragment of MLL (aa3751-3968) expressed in E. Coli.			
Specificity :	MLL Monoclonal Antibody detects endogenous levels of MLL protein.			
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.			
Source :	Monoclonal, Mouse			
Dilution :	WB 1:500 - 1:2000. IHC 1:200 - 1:1000. ELISA: 1:10000 IF 1:50-200			
Purification :	Affinity purification			
Storage Stability -	-15°C to -25°C/1 year(Do not lower than -25°C)			
ctorage otability .				
Molecularweight :	432kD			



P References :	1. Genet Couns. 2006;17(2):155-9. 2. Cancer Genet Cytogenet. 2006 Jul 15;168(2):162-7 2. Laukemia: 2007 Feb;21(2):260 2. Epub 2007 Jap 4		
	5. Leukemia. 2007 Feb,21(2).300-2. Epub 2007 Jan 4.		
Background :	This gene encodes a transcriptional coactivator that plays an essential role in regulating gene expression during early development and hematopoiesis. The encoded protein contains multiple conserved functional domains. One of these domains, the SET domain, is responsible for its histone H3 lysine 4 (H3K4) methyltransferase activity which mediates chromatin modifications associated with epigenetic transcriptional activation. This protein is processed by the enzyme Taspase 1 into two fragments, MLL-C and MLL-N. These fragments reassociate and further assemble into different multiprotein complexes that regulate the transcription of specific target genes, including many of the HOX genes. Multiple chromosomal translocations involving this gene are the cause of certain acute lymphoid leukemias and acute myeloid leukemias. Alternate splicing results in multiple transcript variants.[provided by RefS		
Function :	catalytic activity:S-adenosyl-L-methionine + histone L-lysine = S-adenosyl-L- homocysteine + histone N(6)-methyl-L-lysine.,similarity:Contains 1 post-SET domain.,similarity:Contains 1 SET domain.,		
Subcellular Location :	Nucleus .; [MLL cleavage product N320]: Nucleus.; [MLL cleavage product C180]: Nucleus. Localizes to a diffuse nuclear pattern when not associated with MLL cleavage product N320.		
Expression :	Heart, lung, brain and T- and B-lymphocytes.		

D	м	1	2	-
70 30 52 55			-	
13	-		-	
34		-		←
26				
7				
1				

Products Images

Western Blot analysis using MLL Monoclonal Antibody against truncated MLL recombinant protein (1) and truncated GFP-MLL(aa3714-3969) transfected Cos7 cell lysate (2).





Immunohistochemistry analysis of paraffin-embedded human lung cancer (left) and esophagus cancer (right), showing nuclear localization with DAB staining using MLL Monoclonal Antibody.